



# Routine Vaccine Administration Guide

Please refer to the DHW Publicly Funded Vaccine Schedule [here](#)

# Publicly Funded Vaccines in Nova Scotia

Childhood, Starting in Infancy (By Age)

	VACCINE	PRODUCT NAME	SITE	ROUTE	DOSE	RECONSTITUTION	NEEDLE SIZE	
2 MONTHS	DTaP-IPV-Hib <span style="color:red">!</span>	Pediacel	Vastus lateralis (thigh)	Intramuscular	0.5 mL	No	25 g 1-inch	2 separate needles at least 1-inch apart
	Pneumococcal conjugate (Pneu-C-13)	Prevnar 13	Vastus lateralis (thigh)	Intramuscular	0.5 mL	No		
	Rotavirus	RotaTeq	Orally administered	Oral	2 mL	No	Not applicable	
4 MONTHS	DTaP-IPV-Hib <span style="color:red">!</span>	Pediacel	Vastus lateralis (thigh)	Intramuscular	0.5 mL	No	25 g 1-inch	2 separate needles at least 1-inch apart
	Pneumococcal conjugate (Pneu-C-13)	Prevnar 13	Vastus lateralis (thigh)	Intramuscular	0.5 mL	No		
	Rotavirus	RotaTeq	Orally administered	Oral	2 mL	No	Not applicable	
6 MONTHS	DTaP-IPV-Hib <span style="color:red">!</span>	Pediacel	Vastus lateralis (thigh)	Intramuscular	0.5 mL	No	25 g 1-inch	
	Rotavirus	RotaTeq	Orally administered	Oral	2 mL	No	Not applicable	
12 MONTHS	Pneumococcal conjugate (Pneu-C-13)	Prevnar 13	Vastus lateralis (thigh)	Intramuscular	0.5 mL	No	25 g 1-inch	2 separate needles at least 1-inch apart
	Meningococcal group C conjugate	NeisVac-C Menjugate	Vastus lateralis (thigh)	Intramuscular	0.5 mL	NeisVac-C (No) Menjugate (Yes or pre-filled)		
	MMRV (measles, mumps, rubella, varicella)	Priorix Tetra ProQuad	Upper arm	Subcutaneous	0.5 mL	Yes (vaccine-specific diluent)	25 g 5/8-inch	
18 MONTHS	DTaP-IPV-Hib <span style="color:red">!</span>	Pediacel	Deltoid	Intramuscular	0.5 mL	No	25 g 1-inch	
	MMRV (measles, mumps, rubella, varicella)	Priorix Tetra ProQuad	Upper arm	Subcutaneous	0.5 mL	Yes (vaccine-specific diluent)	25 g 5/8-inch	
4-6 YEARS	Tdap-IPV <span style="color:red">!</span>	Adacel Polio Boostrix Polio	Deltoid	Intramuscular	0.5 mL	No	25 g 1-inch	
ANNUALLY, FROM 6 MONTHS OF AGE ONWARD	Influenza <i>1st-time recipients aged 6 months – 9 years: 2 doses (0, 4 weeks)</i>	Fluzone Flulaval Tetra Afluria (>5 years old only)	6-12 months old: vastus lateralis (thigh) Older than 12 months: deltoid	Intramuscular	0.5 mL	No	25 g 1-inch	

! Vaccines with similar names that may lead to vaccine errors. i Products are subject to change based on availability.



## Catch-up (Resuming or Starting Immunizations)

- Disruptions in vaccine schedules do not require that a series be re-started, regardless of the length of time between doses.
- For catch-up schedules, please refer to the Canadian Immunization Guide for age-specific recommendations:
  - **Less than 7 years old; 7 – 17 years old; 18 years or older**

## Additional Publicly Funded Vaccines for Patients at High-Risk of Vaccine-preventable Disease

Patients at high-risk of vaccine-preventable diseases are eligible for other vaccines based on **condition-specific guidance**.



Updated August 2023



# Routine Vaccine Administration Guide

Please refer to the DHW Publicly Funded Vaccine Schedule [here](#)

## Publicly Funded Vaccines in Nova Scotia

Childhood (By Age)

	PRODUCT NAME	TOTAL DOSES	AGES	SITE	ROUTE	DOSE	RECONSTITUTION	NEEDLE SIZE
<b>DTaP-IPV-Hib</b> !	Pediacel	4	2, 4, 6, 18 months	< 12 months: Vastus lateralis (thigh) Older than 12 months: Deltoid	Intramuscular	0.5 mL	No	25 g 1 inch
<b>ROTAVIRUS</b>	RotaTeq	3	2, 4, 6 months	Orally administered	Oral	2 mL	No	Not applicable
<b>PNEUMOCOCCAL CONJUGATE (PNEU-C-13)</b>	Prenar 13	3	2, 4, 12 months	Vastus lateralis (thigh)	Intramuscular	0.5 mL	No	25 g 1 inch
<b>MENINGOCOCCAL GROUP C CONJUGATE</b>	NeisVac-C Menjugate	1	12 months	Vastus lateralis (thigh)	Intramuscular	0.5 mL	NeisVac-C (No) Menjugate (Yes)	25 g 1 inch
<b>MMRV (MEASLES, MUMPS, RUBELLA, VARICELLA)</b>	Priorix Tetra ProQuad	2	12, 18 months	Upper arm	Subcutaneous	0.5 mL	Yes (vaccine-specific diluent)	25 g 5/8 inch
<b>Tdap-IPV</b> !	Adacel Polio Boostrix Polio	1	4 -6 years	Deltoid	Intramuscular	0.5 mL	No	25 g 1 inch
<b>INFLUENZA</b>	Fluzone Flulaval Tetra Afluria (>5 years old only)	Annually; <i>1st-time recipients aged 6 months – 9 years: 2 doses (0, 4 weeks)</i>	From 6 months of age onward	6-12 months old: vastus lateralis (thigh) Older than 12 months: deltoid	Intramuscular	0.5 mL	No	25 g 1 inch

! Vaccines with similar names that may lead to vaccine errors. i Products are subject to change based on availability.



### Catch-up (Resuming or Starting Immunizations)

- Disruptions in vaccine schedules do not require that a series be re-started, regardless of the length of time between doses.
- For catch-up schedules, please refer to the Canadian Immunization Guide for age-specific recommendations:
  - **Less than 7 years old; 7 – 17 years old; 18 years or older**

### Additional Publicly Funded Vaccines for Patients at High-Risk of Vaccine-preventable Disease

Patients at high-risk of vaccine-preventable diseases are eligible for other vaccines based on **condition-specific guidance**.



Updated August 2023



# Routine Vaccine Administration Guide

Please refer to the DHW Publicly Funded Vaccine Schedule [here](#)

# Publicly Funded Vaccines in Nova Scotia

(Childhood, School, Adult) Starting in Infancy (By Age)

CHILDHOOD VACCINES	VACCINE	PRODUCT NAME	SITE	ROUTE	DOSE	RECONSTITUTION	NEEDLE SIZE
2 MONTHS	DTaP-IPV-Hib <b>!</b>	Pediacel	Vastus lateralis (thigh)	Intramuscular	0.5 mL	No	25 g 1-inch 2 separate needles at least 1-inch apart
	Pneumococcal conjugate (Pneu-C-13)	Prevnar 13	Vastus lateralis (thigh)	Intramuscular	0.5 mL	No	
	Rotavirus	RotaTeq	Orally administered	Oral	2 mL	No	Not applicable
4 MONTHS	DTaP-IPV-Hib <b>!</b>	Pediacel	Vastus lateralis (thigh)	Intramuscular	0.5 mL	No	25 g 1-inch 2 separate needles at least 1-inch apart
	Pneumococcal conjugate (Pneu-C-13)	Prevnar 13	Vastus lateralis (thigh)	Intramuscular	0.5 mL	No	
	Rotavirus	RotaTeq	Orally administered	Oral	2 mL	No	Not applicable
6 MONTHS	DTaP-IPV-Hib <b>!</b>	Pediacel	Vastus lateralis (thigh)	Intramuscular	0.5 mL	No	25 g 1-inch
	Rotavirus	RotaTeq	Orally administered	Oral	2 mL	No	Not applicable
12 MONTHS	Pneumococcal conjugate (Pneu-C-13)	Prevnar 13	Vastus lateralis (thigh)	Intramuscular	0.5 mL	No	25 g 1-inch 2 separate needles at least 1-inch apart
	Meningococcal group C conjugate	Neisvac-C Menjugate	Vastus lateralis (thigh)	Intramuscular	0.5 mL	NeisVac-C (No) Menjugate (Yes)	
	MMRV (measles, mumps, rubella, varicella)	Priorix Tetra ProQuad	Upper arm	Subcutaneous	0.5 mL	Yes (vaccine-specific diluent)	25 g 5/8-inch
18 MONTHS	DTaP-IPV-Hib <b>!</b>	Pediacel	Deltoid	Intramuscular	0.5 mL	No	25 g 1-inch
	MMRV (measles, mumps, rubella, vari-cella)	Priorix Tetra ProQuad	Upper arm	Subcutaneous	0.5 mL	Yes (vaccine-specific diluent)	25 g 5/8-inch
4-6 YEARS	Tdap-IPV <b>!</b>	Adacel Polio Boostrix Polio	Deltoid	Intramuscular	0.5 mL	No	25 g 1-inch

## SCHOOL VACCINES - Catch-up vaccines available through Public Health and/or Primary Care Provider prior to 19th Birthday

GRADE 7 SCHOOL VACCINES	HPV series (0, 6 months)	Gardasil 9	Deltoid	Intramuscular	0.5 mL	No	25 g 1-inch
	Hepatitis B series (0, 6 months)	Engerix-B Recombivax	Deltoid	Intramuscular	1.0 mL	No	25 g 1-inch
	Tdap	Adacel Boostrix	Deltoid	Intramuscular	0.5 mL	No	25 g 1-inch
	Meningococcal Quadrivalent (ACYW135)	Menveo Menactra Nimenrix	Deltoid	Intramuscular	0.5 mL	No	25 g 1-inch

**!** Vaccines with similar names that may lead to vaccine errors. **i** Products are subject to change based on availability.



# Routine Vaccine Administration Guide

Please refer to the DHW Publicly Funded Vaccine Schedule [here](#)

# Publicly Funded Vaccines in Nova Scotia

(Childhood, School, Adult) Starting in Infancy (By Age)

ADULT VACCINES	VACCINE	PRODUCT NAME	SITE	ROUTE	DOSE	RECONSTITUTION	NEEDLE SIZE
EVERY 10 YEARS	Td (Booster)	Td adsorbed	Deltoid	Intramuscular	0.5 mL	No	25 g 1-inch
ONCE AS AN ADULT	Tdap <i>*If not previously given in adulthood. May receive one dose 10 years after receiving a tetanus containing vaccine.</i>	Adacel Boostrix	Deltoid	Intramuscular	0.5 mL	No	25 g 1-inch
PREGNANT PERSONS	Tdap: offered every pregnancy between 27 and 32 weeks	Adacel Boostrix	Deltoid	Intramuscular	0.5 mL	No	25 g 1-inch
ADULTS BORN IN 1970 OR LATER	MMR (Booster)	MMRII (SC only) Priorix (SC preferred)	SC: Upper Arm IM: Deltoid	Subcutaneous Intramuscular	0.5 mL	Yes (vaccine-specific diluent)	SC: 25g 5/8-inch IM: 25g 1-inch
<b>VACCINES FOR SENIORS</b>							
65 YEARS AND OLDER	Pneumococcal polysaccharide (Pneu-P-23)	Pneumovax 23	IM: Deltoid SC: Upper Arm	Intramuscular Subcutaneous	0.5 mL	No	IM: 25 g 1-inch SC: 25 g 5/8-inch
65 YEARS AND OLDER	High-dose influenza; annually	Fluzone HD	Deltoid	Intramuscular	0.7 mL	No	25 g 1-inch
<b>VACCINES FOR ALL AGES</b>							
ANNUALLY, FROM 6 MONTHS OF AGE ONWARD	Influenza <i>1st-time recipients aged 6 months – 9 years: 2 doses (0, 4 weeks)</i>	Fluzone Quad Flulaval Tetra Afluria Tetra (>5 years old only)	6-12 months old: vastus lateralis (thigh) Older than 12 months: deltoid	Intramuscular	0.5 mL	No	25 g 1-inch

**i** For information on vaccination and pregnancy see [here](#). **i** For information on Covid-19 vaccination see [here](#). **i** Products are subject to change based on availability.



## Catch-up (Resuming or Starting Immunizations)

- Disruptions in vaccine schedules do not require that a series be re-started, regardless of the length of time between doses.
- For catch-up schedules, please refer to the Canadian Immunization Guide for age-specific recommendations:
  - **Less than 7 years old; 7 – 17 years old; 18 years or older**

## Additional Publicly Funded Vaccines for Patients at High-Risk of Vaccine-preventable Disease

Patients at high-risk of vaccine-preventable diseases are eligible for other vaccines based on **condition-specific guidance**.



Updated August 2023